



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

The relationship between *Helicobacter pylori*-related microbiota dysbiosis and gastrointestinal tract pathologies

Kountouras, Jannis ; Doulberis, Michael ; Papaefthymiou, Apostolis ; Polyzos, Stergios A ; Toulountzi, Maria ; Vardaka, Elisabeth ; Kapetanakis, Nikolaos ; Liatsos, Christos ; Gavalas, Emmanouel ; Anastasiadis, Sotiris ; Tzivras, Dimitri ; Katsinelos, Panagiotis

DOI: <https://doi.org/10.1080/00365521.2019.1619826>

Posted at the Zurich Open Repository and Archive, University of Zurich
ZORA URL: <https://doi.org/10.5167/uzh-181170>
Journal Article

Originally published at:

Kountouras, Jannis; Doulberis, Michael; Papaefthymiou, Apostolis; Polyzos, Stergios A; Toulountzi, Maria; Vardaka, Elisabeth; Kapetanakis, Nikolaos; Liatsos, Christos; Gavalas, Emmanouel; Anastasiadis, Sotiris; Tzivras, Dimitri; Katsinelos, Panagiotis (2019). The relationship between *Helicobacter pylori*-related microbiota dysbiosis and gastrointestinal tract pathologies. *Scandinavian Journal of Gastroenterology*, 54(6):806-807.

DOI: <https://doi.org/10.1080/00365521.2019.1619826>

The relationship between *Helicobacter pylori*-related microbiota dysbiosis and gastrointestinal tract pathologies

**Jannis Kountouras, MD, PhD¹, Michael Doulberis, MD,DVM, PhD^{1,2},
Apostolis Papaefthimiou, MD^{1,3}, Stergios A. Polyzos, MD, Msc, PhD^{1,4},
Maria Touloumtzi, MD¹, Elisabeth Vardaka, PhD¹, Nikolaos Kapetanakis,
MD, PhD¹, Christos Liatsos, MD, PhD^{1,3}, Emmanouel Gavalas, MD, PhD¹,
Sotiris Anastasiadis, MD¹, Dimitri Tzivras, MD, PhD⁵, Panagiotis
Katsinelos, MD, PhD¹.**

¹Department of Gastroenterology, Second Medical Clinic, Aristotle University of Thessaloniki, Ippokration Hospital, Thessaloniki, Greece

²Department of Gastroenterology and Hepatology, University of Zurich, Zurich 8091, Switzerland

³Department of Gastroenterology, 401 General Military Hospital of Athens, Athens 11525, Greece

⁴First Department of Pharmacology, Medical School, Aristotle University of Thessaloniki, Thessaloniki 54642, Macedonia, Greece

⁵St. Elisabeth-Hospital Herten GmbH, 45699 Herten, Germany

Key words: *Helicobacter pylori*; gastric microbiota; dysbiosis; gastric cancer; esophageal cancer

Conflict of Interest: None declared

Please address correspondence to:

Jannis Kountouras, MD, PhD

Professor of Medicine

8 Fanariou St, Byzantio

551 33, Thessaloniki, Macedonia,

Greece

Tel: +30-2310-892238, Fax: +30-2310-992794

E-mail: jannis@auth.gr, ancorarus2010@gmail.com

To the Editor,

In their review, Zhang et al. [1], concluded that gastrointestinal microbiota (GI-M) plays an essential role in the development and progression of gastric pathologies; they considered a potential impact of *Helicobacter pylori* infection (*Hp*-I) on gastric microbiota (GM) composition in gastric cancer (GC) pathophysiology, though it remains to be elucidated.

In this regard, apart from the mentioned gastric microbiota influenced by *Hp* colonization, the authors did not take into consideration specific pathogens including species from *Campylobacter* genus that appear to be involved in gastric carcinogenesis. *Hp* abundance is positively correlated with the presence of *Campylobacter* which is the most well-connected and influential bacterium observed in *Hp*-related atrophic gastritis patients and could be a substantial contributor to gastric carcinogenesis [2,3].

In this concern, it has been proposed that chronic *Hp*-I induces atrophic gastritis accompanying by decreased acid secretion and acid reflux, thereby reducing the risk of gastroesophageal reflux disease (GERD) and its complications Barrett's esophagus (BE) and esophageal adenocarcinoma (EAC) [4]. However, this conventional consideration might represent a double-edged sword one view. Regarding the opposed view, hypochlorhydria induced by *Hp*-related atrophic gastritis, results in dysbiosis of GI-M, which, beyond GC, could also contribute to BE-EAC sequence [3,5]. Data on BE biofilm show high level of atypical nitrate reducing *Campylobacter* species in BE, compared with non-BE specimens [6] and these species might contribute to development, exacerbation and/or BE progression to EAC through chronic inflammation [5]. Therefore, *Campylobacter* species and other pathogens-related dysbiosis associated with *Hp*-induced atrophic gastritis might be etiological agents of the chronic esophageal inflammation leading to EAC development [5,7] and thus further investigation is needed.

The authors considered molecular mechanisms involved in *Hp*-I related gastric

cancerogenesis [1]. Likewise, concerning the role of molecular events involved in *Hp*-I-related GERD- BE-EAC sequence, we recently summarized [8] comparable *Hp*-related mechanisms such as the following: *Hp* induction of upper and lower gastrointestinal tract (GIT) oncogenic gastrin, which stimulates proliferation via Janus Kinase (JAK) 2 and Akt-dependent nuclear factor-kappa B (NF- κ B) activation in Barrett's EAC cells, displays a anti-apoptotic effect through Bcl-2 protein and survivin upregulations, and induces the mitogenic and oncogenic cyclooxygenase (COX)-2 expression. *Hp* activates the mentioned NF- κ B, a transcription regulator of inflammatory genes, including COX-2 that regulates gastrointestinal neoplasm cell growth and proliferation. Prostaglandins (PGs) derived from upregulated COX-2 contribute to BE cancer progression, by perpetuating chronic inflammation and the mitogenic and antiapoptotic effects of PGs are mediated through activation of several aforementioned signaling pathways including NF- κ B, Src, JAK2/STAT3, ERK, MAPK and PI3K/Akt kinases. Moreover, *Hp*-I could induce specific molecular changes (genetic instability, E-cadherin methylation, monoclonal antibody Das-1) linked with BE pathophysiology, and promotes Ki-67 expression predicting malignant progression in BE. Finally, *Hp*-related metabolic syndrome (MetS) disorders that associated with GI-M dysbiosis appear to be involved in gastric and esophageal carcinogenesis [9,10]. Again, we recently reviewed several *Hp*-related MetS mechanisms involved in upper and lower GIT oncogenic processes [9].

Therefore, *Hp* eradication and manipulation of its related GI-M dysbiosis might inhibit the aforementioned GIT oncogenic processes, and thus further studies are necessary.

References

- [1] Zhang S, Shi D, Li M, et al. The relationship between gastric microbiota and gastric disease. Scand J Gastroenterol. 2019 Apr 4:1-6.
- [2] Thorell K, Bengtsson-Palme J, Liu OH, et al. *In Vivo* Analysis of the Viable Microbiota and Helicobacter pylori Transcriptome in Gastric Infection and Early

Stages of Carcinogenesis. *Infect Immun*. 2017;85(10). pii: e00031-17.

- [3] Parsons BN, Ijaz UZ, D'Amore R, et al. Comparison of the human gastric microbiota in hypochlorhydric states arising as a result of *Helicobacter pylori*-induced atrophic gastritis, autoimmune atrophic gastritis and proton pump inhibitor use. *PLoS Pathog*. 2017;13: e1006653.
- [4] Tomasello G, Giordano F, Mazzola M, et al. *Helicobacter pylori* and Barrett's esophagus: a protective factor or a real cause? *J Biol Regul Homeost Agents*. 2017;31:9-15.
- [5] Mozaffari Namin B, Soltan Dallal MM, Ebrahimi Daryani N. The Effect of *Campylobacter concisus* on Expression of IL-18, TNF- α and p53 in Barrett's Cell Lines. *Jundishapur J Microbiol*. 2015; 8:e26393.
- [6] Blackett KL, Siddhi SS, Cleary S, et al. Oesophageal bacterial biofilm changes in gastro-oesophageal reflux disease, Barrett's and oesophageal carcinoma: association or causality? *Aliment Pharmacol Ther*. 2013; 37: 1084–1092.
- [7] Yang L, Chaudhary N, Baghdadi J, et al. Microbiome in reflux disorders and esophageal adenocarcinoma. *Cancer J*. 2014;20:207-210.
- [8] Kountouras J, Polyzos SA, Doulberis M, et al. Potential impact of *Helicobacter pylori*-related metabolic syndrome on upper and lower gastrointestinal tract oncogenesis. *Metabolism*. 2018;87:18-24.
- [9] Kountouras J, Boziki M, Polyzos SA, et al. The Emerging Role of *Helicobacter Pylori*-Induced Metabolic Gastrointestinal Dysmotility and Neurodegeneration. *Curr Mol Med*. 2017;17:389-404.
- [10] Rogers CJ, Prabhu KS, Vijay-Kumar M. The microbiome and obesity-an established risk for certain types of cancer. *Cancer J*. 2014;20:176-180.